



PCT/GB 2003 / 005179

Rec'd PCT/PTO 03 JUN 2003
10/537346

INVESTOR IN PEOPLE

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

REC'D 05 FEB 2004

WIPO PCT


I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

I also certify that the application is now proceeding in the name as identified herein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.



Signed *AmBrewster*

Dated 22 January 2004

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY



INVESTOR IN PEOPLE

GB 0228365.3

By virtue of a direction given under Section 30 of the Patents Act 1977, the application is proceeding in the name of:

VERNALIS (OXFORD) LIMITED,
Watlington Road,
OXFORD,
OX4 5LY,
United Kingdom

Incorporated in the United Kingdom,

[ADP No. 08740102001]

Patents Form 1/77

Patents Act 1977

Rules 16

THE PATENT OFFICE
A
Office

- 5 DEC 2002

1/77

Request for grant of a patent

The Patent Office

Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference	241/AJW		
2. Patent application number	0228365.3		
3. Full name, address and postcode of the or of each applicant	British Biotech Pharmaceuticals Ltd Watlington Road Cowley Oxford OX4 5LY Patents ADP number 937800 If the applicant is a corporate body, give the country/state of its corporation GB		
4. Title of invention	Antibacterial Agents		
5. Name of your agent	Alan J. Walls		
Address for service in the United Kingdom to which all correspondence should be sent	British Biotech Pharmaceuticals Ltd Watlington Road Cowley Oxford OX4 5LY		
Patents ADP number	17442004		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application no.	Date of filing (day/month/year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier applications		Date of filing (day/month/year)
8. Is a statement of inventorship and of right to grant of a patent required in support of this request?	Yes		

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form	0
Description	30 /
Claims(s)	8 /
Abstract	0
Drawing(s)	0

CF

10. If you are also filing any of the following, state how many against each item.

Priority documents	0
Translations of priority documents	0
Statement of inventorship and right to grant a patent	0
Request for preliminary examination	0
Request for substantive examination	0
Any other documents	0

11. I/We request the grant of a patent on the basis of this application.

Signature

A. J. Walls

Date

04/12/2002

Anthony Weir
Director

For and on behalf of British Biotech Pharmaceuticals Ltd

12. Name and daytime telephone number of person to contact in the United Kingdom

Alan J. Walls
01865 748747

Antibacterial Agents

This invention relates to novel hydroxamic acid and N-formyl hydroxylamine derivatives having antibacterial activity, to methods of treatment using such compounds, and to pharmaceutical and veterinary compositions comprising such compounds.

Background to the Invention

Many classes of antibacterial agents are known, including the penicillins and cephalosporins, tetracyclines, sulfonamides, monobactams, fluoroquinolones and quinolones, aminoglycosides, glycopeptides, macrolides, polymyxins, lincosamides, trimethoprim and chloramphenicol. The fundamental mechanisms of action of these antibacterial classes vary.

Bacterial resistance to many known antibacterials is a growing problem. Accordingly there is a continuing need in the art for alternative antibacterial agents, especially those which have mechanisms of action fundamentally different from the known classes, and/or which are effective against the causative organisms of community acquired respiratory infections, and/or which are selective in their pharmacological activity, thus reducing risk of unwanted side effects..

Amongst the Gram-positive pathogens, such as staphylococci, streptococci, mycobacteria and enterococci, resistant strains have evolved/arisen which makes them particularly difficult to eradicate. Examples of such strains are methicillin resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase negative Staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* and multiply resistant *Enterococcus faecium*.

Brief Description of the Invention

This invention makes available a new class of hydroxamic acid and N-formyl hydroxylamine derivatives having antibacterial activity. The compounds are characterised inter alia by the presence of a cyclic diazole feature in their structural

backbone.

Although it may be of interest to establish the mechanism of action of the compounds with which the invention is concerned, it is their ability to inhibit bacterial growth that makes them useful. However, it is presently believed that their antibacterial activity is due, at least in part, to intracellular inhibition of bacterial polypeptide deformylase (PDF; EC 3.5.1.31).

Related Prior Art


Although there are many publications disclosing both hydroxamic acid and N-formyl hydroxylamine derivatives as inhibitors of various metalloenzymes such as angiotensin converting enzyme, enkephalinase and the matrix metalloproteinases, there are relatively few relating to such compounds as antibacterial agents. The following patent publications are relevant in that connection:

WO 99/39704	(British Biotech)
WO 99/57097	(Versicor)
WO 99/59568	(British Biotech)
WO 00/35440	(British Biotech)
WO 00/44373	(British Biotech)
WO 00/58294	(British Biotech)
WO 00/61134	(British Biotech)
WO 01/10835	(British Biotech)
WO 01/38561	(Questcor)
WO 01/40198	(Aventis)
WO 01/42431	(Bayer)
WO 01/44178	(Versicor)
WO 01/44179	(Versicor)
WO 01/85160	(SmithKline Beecham)
WO 01/85170	(SmithKline Beecham)
WO 02/28829	(Questcor)
WO 02/41886	(British Biotech)

WO 02/50081 (British Biotech)


WO 02/070541 (SmithKline Beecham) relates to PDF inhibitor compounds which have a hydrazide feature in the backbone and an N-formylhydroxylamino metal binding group.

In addition the natural antibiotic actinonin (see for example J.C.S Perkin I, 1975, 819) is a hydroxamic acid derivative of Structure (A):



which is now known to act by inhibition of PDF. In addition to actinonin, various structural analogues of actinonin have also been shown to have antibacterial activity (see for example Broughton et al. (Devlin et al. Journal of the Chemical Society. Perkin Transactions 1 (9):830-841, 1975; Broughton et al. Journal of the Chemical Society. Perkin Transactions 1 (9):857-860, 1975).

The matlystatin group of compounds share a number of structural similarities with actinonin. Both are peptidic molecules with functional hydroxamic acid metal binding groups (Ogita et al., J. Antibiotics. 45(11):1723-1732; Tanzawa et al., J. Antibiotics. 45(11):1733-1737; Haruyama et al., J. Antibiotics. 47(12):1473-1480; Tamaki et al., J. Antibiotics. 47(12):1481-1492). The matlystatins and their close structural analogues are characterised by the presence in the molecule of a divalent piperazin-1, 6-diyl group, i.e.



In view of their close structural similarity to actinonin, the observation that actinonin inhibits PDF implies that matlystatin compounds may also inhibit PDF.

For a recent review of peptide deformylase inhibitors, see Clements et. al., Curr. Med. Chem. - Anti-Infective Agents, 2002, 1, 239-249.

The following patent publications disclose N-formyl hydroxylamine structures:

EP-B-0236872	(Roche)
WO 92/09563	(Glycomed)
WO 92/04735	(Syntex)
WO 95/19965	(Glycomed)
WO 95/22966	(Sanofi Winthrop)
WO 95/33709	(Roche)
WO 96/23791	(Syntex)
WO 96/16027	(Syntex/Agouron)
WO 97/03783	(British Biotech)
WO 97/18207	(DuPont Merck)
WO 98/38179	(GlaxoWellcome)
WO 98/47863	(Labs Jaques Logeais)


The pharmaceutical utility ascribed to the N-formyl hydroxylamine derivatives in those publications is the ability to inhibit matrix metalloproteinases (MMPs) and in some cases release of tumour necrosis factor (TNF), and hence the treatment of diseases or conditions mediated by those enzymes, such as cancer and rheumatoid arthritis.

In addition to these, US-A-4,738,803 (Roques et al.) also discloses N-formyl hydroxylamine derivatives. However, these compounds are disclosed as enkephalinase inhibitors and are proposed for use as antidepressants and hypotensive agents. Also, WO 97/38705 (Bristol-Myers Squibb) discloses certain N-formyl hydroxylamine derivatives as enkephalinase and angiotensin converting enzyme inhibitors.

There are too many publications relating to metalloenzyme inhibiting hydroxylamic acid derivatives to summarise effectively. However a recent review, Whittaker et. al. Chem. Rev. 1999, 99, 2735, provides an overview of that art.

Description of the invention

The present invention provides a compound of formula (I) or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof



wherein

Q represents a radical of formula $-\text{N}(\text{OH})\text{CH}(=\text{O})$ or formula $-\text{C}(=\text{O})\text{NH}(\text{OH})$;

Y represents $-\text{C}(=\text{O})-$, $-\text{C}(=\text{S})-$, $-\text{S}(=\text{O})-$, or $-\text{SO}_2-$;

R_1 represents hydrogen, $\text{C}_1\text{-C}_6$ alkyl or $\text{C}_1\text{-C}_6$ alkyl substituted by one or more halogen atoms, or, except when Q is a radical of formula $-\text{N}(\text{OH})\text{CH}(=\text{O})$, a hydroxy, $\text{C}_1\text{-C}_6$ alkoxy, $\text{C}_1\text{-C}_6$ alkenyloxy, halogen, amino, $\text{C}_1\text{-C}_6$ alkylamino, or di- ($\text{C}_1\text{-C}_6$ alkyl)amino group;

R_2 represents a substituted or unsubstituted $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_1\text{-C}_3$ alkyl-O- $\text{C}_1\text{-C}_3$ alkyl,

C₁-C₃ alkyl-S-C₁-C₃ alkyl, cycloalkyl(C₁-C₃ alkyl)-, aryl(C₁-C₃ alkyl)-, heterocyclyl(C₁-C₃ alkyl)-, or R¹R²N-C₁-C₃ alkyl group wherein R¹ represents hydrogen or C₁-C₃ alkyl and R² represents C₁-C₃ alkyl, or R¹R²N- represents a cyclic amino group;

R₃ and R₄ taken together with the nitrogen atoms to which they are respectively attached form a saturated heterocyclic ring of from 4 to 7 ring atoms, which may be fused to a second carbocyclic or heterocyclic ring, either of which rings may optionally be substituted; and

A represents a primary, secondary or tertiary amino group or a group -R₅, -OR₅, wherein R₅ is a substituted or unsubstituted C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, cycloalkyl, aryl, heterocyclyl, C₁-C₃ alkyl-O-C₁-C₃ alkyl, C₁-C₃ alkyl-S-C₁-C₃ alkyl, cycloalkyl(C₁-C₃ alkyl)-, heterocyclic(C₁-C₃ alkyl, aryl(C₁-C₃ alkyl)-, or R¹R²N-C₁-C₃ alkyl group wherein R¹ represents hydrogen or C₁-C₃ alkyl and R² represents C₁-C₃ alkyl, or R¹R²N- represents a cyclic amino group.

In another aspect, the invention provides a method for the treatment of bacterial infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially effective dose of a compound of formula (I) as defined above. Also included in the invention is the use of a compound of formula (I) as defined above for inhibiting bacterial growth in vitro and in vivo in mammals, and the use of such a compound for the manufacture of a composition for treating bacterial infection by inhibiting bacterial growth.

In a further aspect of the invention there is provided a method for the treatment of bacterial contamination by applying an antibacterially effective amount of a compound of formula (I) as defined above to the site of contamination.

The compounds of formula (I) as defined above may be used as component(s) of antibacterial cleaning or disinfecting materials.

As used herein terms of the form "(C_a-C_b)alkyl" where a and b are integers refer to a

straight or branched chain alkyl moiety having from a to b carbon atoms. Thus, for example, the term "(C₁-C₆)alkyl" means a straight or branched chain alkyl moiety having from 1 to 6 carbon atoms, and includes, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, t-butyl, n-pentyl and n-hexyl.

As used herein terms of the form "divalent (C_a-C_b)alkylene radical" where a and b are integers refer to a saturated hydrocarbon chain having from a to b carbon atoms and two unsatisfied valencies.

As used herein terms of the form "(C_a-C_b)alkenyl" where a and b are integers refer to straight or branched chain alkenyl moiety having from a to b carbon atoms having at least one double bond of either E or Z stereochemistry where applicable. Thus, for example, the term "(C₁-C₆)alkenyl" means a straight or chain alkenyl moiety having from 2 to 6 carbon atoms having at least one double bond, and includes, for example, vinyl, allyl, 1- and 2-butenyl and 2-methyl-2-propenyl.

As used herein the term "C_a-C_b alkynyl" where a and b are integers refers to straight chain or branched chain hydrocarbon groups having from two to six carbon atoms and having in addition one triple bond. Thus, for example, the term "(C₁-C₆)alkynyl" would include for example, ethynyl, 1-propynyl, 1- and 2-butyne, 2-methyl-2-propynyl, 2-pentyne, 3-pentyne, 4-pentyne, 2-hexynyl, 3-hexynyl, 4-hexynyl and 5-hexynyl.

As used herein the term "cycloalkyl" means a saturated alicyclic moiety having from 3-8 carbon atoms and includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and cyclooctyl.

As used herein the term "aryl" refers to a mono-, bi- or tri-cyclic carbocyclic aromatic group, and to groups consisting of two covalently linked monocyclic carbocyclic aromatic groups. Illustrative of such groups are phenyl, biphenyl and naphthyl.

As used herein the term "heteroaryl" refers to a 5- or 6- membered aromatic ring

containing one or more heteroatoms. Illustrative of such groups are thienyl, furyl, pyrrolyl, imidazolyl, benzimidazolyl, thiazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl.

As used herein the unqualified term "heterocyclyl" or "heterocyclic" includes "heteroaryl" as defined above, and in particular means a 5-7 membered aromatic or non-aromatic heterocyclic ring containing one or more heteroatoms selected from S, N and O, including for example, pyrrolyl, furanyl, thienyl, piperidinyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, thiadiazolyl, pyrazolyl, pyridinyl, pyrrolidinyl, pyrimidinyl, morpholinyl, piperazinyl, indolyl, morpholinyl, benzofuranyl, pyranyl, isoxazolyl, benzimidazolyl, methylenedioxyphenyl, maleimido and succinimido groups.

Unless otherwise specified in the context in which it occurs, the term "substituted" as applied to any moiety herein means substituted with up to four substituents, each of which independently may be (C₁-C₆)alkyl, (C₁-C₆)alkoxy, hydroxy, mercapto, (C₁-C₆)alkylthio, amino, halo (including fluoro, chloro, bromo and iodo), trifluoromethyl, nitro, -COOH, -CONH₂, -COR^A, -COOR^A, -NHCOR^A, -CONHR^A, -NHR^A, -NR^AR^B, or -CONR^AR^B wherein R^A and R^B are independently a (C₁-C₆)alkyl group.

Salts of the compounds of the invention include physiologically acceptable acid addition salts for example hydrochlorides, hydrobromides, sulphates, methane sulphonates, p-toluenesulphonates, phosphates, acetates, citrates, succinates, lactates, tartrates, fumarates and maleates. Salts may also be formed with bases, for example sodium, potassium, magnesium, and calcium salts.

There are several actual or potential chiral centres in the compounds according to the invention because of the presence of asymmetric carbon atoms. The presence of several asymmetric carbon atoms gives rise to a number of diastereoisomers with R or S stereochemistry at each chiral centre. The invention includes all such diastereoisomers and mixtures thereof. Currently, the preferred stereoconfiguration

of the carbon atom carrying the R_2 group is R.

In the compounds of the invention, in relation to the groups Q, R_1 , R_2 , R_3 , R_4 , Y and A, separately and in combination:

The group Q

It is currently preferred that Q is an N-formyl hydroxylamine group $-N(OH)CH(=O)-$.

The radical -Y-

It is currently preferred that -Y- is $-C(=O)-$ or $-SO_2-$.

The group R_1

R_1 may be, for example, hydrogen, methyl, trifluoromethyl or, in the case where Q is a hydroxamic acid group $HONHCO-$, fluorine. Hydrogen is currently preferred.

The group R_2

R_2 may be, for example:

optionally substituted C_1-C_6 alkyl, C_3-C_6 alkenyl, C_3-C_6 alkynyl or cycloalkyl;

phenyl(C_1-C_6 alkyl)-, phenyl(C_3-C_6 alkenyl)- or phenyl(C_3-C_6 alkynyl)-
optionally substituted in the phenyl ring;

cycloalkyl(C_1-C_6 alkyl)-, cycloalkyl(C_3-C_6 alkenyl)- or cycloalkyl(C_3-C_6 alkynyl)-
optionally substituted in the cycloalkyl ring; or

$CH_3(CH_2)_pO(CH_2)_q-$ or $CH_3(CH_2)_pS(CH_2)_q-$, wherein p is 0, 1, 2 or 3 and q is 1, 2 or 3.

Specific examples of R_2 groups include


methyl, ethyl, n- and iso-propyl, n- and iso-butyl, n-pentyl, iso-pentyl, 3-methyl-but-1-yl, n-hexyl, n-heptyl, n-acetyl, n-octyl, methylsulfanylethyl, ethylsulfanylmethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-ethoxymethyl, 3-

hydroxypropyl, allyl, 3-phenylprop-3-en-1-yl, prop-2-yn-1-yl, 3-phenylprop-2-yn-1-yl, 3-(2-chlorophenyl)prop-2-yn-1-yl, but-2-yn-1-yl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl, cyclohexylmethyl, cyclohexylethyl, cyclohexylpropyl, furan-2-ylmethyl, furan-3-methyl, tetrahydrofuran-2-ylmethyl, tetrahydrofuran-2-ylmethyl, piperidinylmethyl, pyrid-2-ylmethyl, pyrid-3-ylmethyl, pyrid-4-ylmethyl, phenylpropyl, 4-chlorophenylpropyl, 4-methylphenylpropyl, 4-methoxyphenylpropyl, benzyl, 4-chlorobenzyl, 4-methylbenzyl, and 4-methoxybenzyl.

Presently preferred groups at R_2 are (C_1-C_6) alkyl-, cycloalkylmethyl-, (C_1-C_3) alkyl-S- (C_1-C_3) alkyl-, or (C_1-C_3) alkyl-O- (C_1-C_3) alkyl-, especially n-propyl, n-butyl, n-pentyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl or cyclohexylethyl.

The ring formed by R_3 and R_4 and the nitrogens to which they are attached

Examples of such rings are the following, wherein R represents hydrogen or C_1-C_4 alkyl




In the above rings, where a sulfur atom is present as a ring member, the equivalent structures wherein that sulfur is oxidised to $-SO-$ or $-SO_2-$ are also examples of ring structures which may be formed by R_3 and R_4 and the nitrogens to which they are attached.

The group A

The group A is a primary, secondary or tertiary amino group or a group $-R_5$, or $-OR_5$. When A is $-R_5$, or $-OR_5$, the R_5 group may be, for example, any of those given as R_2 examples above, or a group of formula (II) as defined below, including such specific

examples of groups (II) as morpholinyl, furanyl, thienyl, phenyl, and benzyl..

Presently it is preferred that A is a secondary or tertiary amino group, and in the latter case it may be a non-cyclic or a cyclic amino group. For example, A may be an amino group of formula -NR₆R₇ wherein R₆ and R₇ independently represent a radical of formula (II)



wherein

m, p and n are independently 0 or 1;

Z represents hydrogen or a carbocyclic or heterocyclic ring of 5 to 7 ring atoms which is optionally fused to a saturated or unsaturated carbocyclic or heterocyclic second ring of 5 to 7 ring atoms;

Alk¹ and Alk² independently represent divalent C₁-C₃ alkylene radicals;

X represents -O-, -S-, -S(O)-, -S(O₂)-, -C(=O)-, -NH-, -NR₇- where R₇ is C₁-C₃ alkyl;

and wherein

Alk¹, Alk² and Z when other than hydrogen, independently are optionally substituted by

(C₁-C₃)alkyl, (C₂-C₃)alkenyl, or (C₂-C₃)alkynyl,
phenyl, optionally substituted by (C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo, nitro,
amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl;

monocyclic 5 or 6-membered heterocyclic, optionally substituted by (C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo, nitro, amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl
benzyl, optionally substituted in the phenyl ring by (C₁-C₃)alkyl, (C₁-C₃)alkoxy,

halo, nitro, amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl,
 hydroxy, phenoxy, (C₁-C₆)alkoxy, or hydroxy(C₁-C₆)alkyl,
 mercapto, (C₁-C₆)alkylthio or mercapto(C₁-C₆)alkyl,
 oxo,
 nitro,
 cyano
 halo
 -COOH, or -COOR^A,
 -CONH₂, -CONHR^A, or -CONR^AR^B
 -COR^A, -SO₂R^A,
 -NHCOR^A,
 -NH₂, -NHR^A, or -NR^AR^B,

wherein R^A and R^B are independently a (C₁-C₆) alkyl group, R^A and R^B taken together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclic ring which may be substituted by (C₁-C₃)alkyl, hydroxy, or hydroxy(C₁-C₃)alkyl.

The amino group A may also be one of formula -NR₆R₇ wherein R₆ and R₇ when taken together with the nitrogen atom to which they are attached form a saturated heterocyclic ring of 5 to 8 atoms optionally fused to a saturated or unsaturated carbocyclic or heterocyclic second ring of 5 to 7 ring atoms, any of which rings being optionally substituted by a radical of formula (II) as defined above. Examples of cyclic amino groups are 1-pyrrolidinyl, piperidin-1-yl, 1-piperazinyl, hexahydro-1-pyridazinyl, morpholin-4-yl, tetrahydro-1,4-thiazin-4-yl, tetrahydro-1,4-thiazin-4-yl 1-oxide, tetrahydro-1,4-thiazin-4-yl 1,1-dioxide, hexahydroazipino, thiomorpholino, diazepino, and thiazolidinyl. Presently preferred are piperidin-1-yl and 1-piperazinyl.

The group of formula (II)


In the substituent (II) Alk¹ and Alk² may independently represent, for example

$-(CH_2)-$ or $-(CH_2CH_2)-$. In the case where m is 0 and p is 1, X may be, for example $-S-$, $-S(=O)-$, or preferably $-C(=O)-$ or $-S(O_2)-$. In such cases n may be 0 or 1, and when A is a cyclic amino group which contains a second ring nitrogen, the X radical may be linked to that ring nitrogen, for example (in the case of $X = -C(=O)-$ or $-S(O_2)-$) as an amide or sulphonamide bond.


In the substituent (II) m , n and p may all be 0, so that the group Z is directly linked to the amino group A .

In a preferred subset of the compounds of the invention, the substituent (II) has the formula $-CH_2Z$, $-OZ$, or $-(C=O)Z$ wherein Z is C_1 - C_3 alkyl, phenyl, 3,4-methylenedioxyphenyl, morpholinyl, pyrimidinyl, 1,2,3-thiadiazolyl, 1,4-thiazolyl, benzofuranyl, furanyl, thienyl, pyranal, pyrrolyl, pyrazolyl, isoxazolyl, or pyridyl, any of which may optionally be substituted as specified. In particular, Z may be a methyl, ethyl, n - or iso-propyl, phenyl, 3,4-methylenedioxyphenyl, morpholinyl, pyrimidin-2-yl, 1,2,3-thiadiazol-5-yl, 1,4-thiazol-5-yl, benzofuran-2-yl, 2- or 3-furanyl, 2- or 3-thienyl, 2- or 3-pyranal, 2-, 3- or 4-pyrrolyl, 3-, 4- or 5-pyrazolyl, 3-, 4- or 5-isoxazolyl, or 2-, 3- or 4-pyridyl ring any of which may optionally be substituted as specified in the broad description of the compounds of the invention.

Compounds of the invention include those selected from the group consisting of compounds of formulae (IIA) or (IIB).



(IIA)



(IIB)



wherein R_2 is as defined in relation to formula (II), especially n -propyl, n -butyl, n -pentyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl or cyclohexylethyl;

X_1 is a bond, C_1-C_3 alkylene, $-NH-$ or $-O-$; and




A_1 is optionally substituted C_1-C_6 alkyl, cycloalkyl, aryl, or heterocyclic, for example methyl, ethyl phenyl, cyclopentyl, cyclohexyl, 2- or 3-furanyl, 2- or 3-thienyl, 2-, 3- or 4-pyridyl, 3-, 4- or 5-pyrazolyl, 3-, 4- or 5-oxazolyl, or 3-, 4- or 5-thiazolyl, methoxymethyl, 3,5-bis-(trifluoromethyl)phenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4-methylenedioxyphenyl, 4-fluorophenyl benzyl, 3-pyridyl, 4-pyridyl, cyclohexyl, 1,3-dimethylpyrazol-5-yl, 1-methylimidazol-5-yl, and 2-[morpholin-1-yl]pyrid-5-yl.

Particular compounds of the invention include those of the Examples herein.

In general, compounds of the invention are accessible by conventional synthetic procedures, for example by acylation of cyclic diazole compound (IV) with an activated derivative of an acid (III),



by methods known from peptide synthesis for example using an acid chloride, wherein the NH and/or OH groups in Q are protected during the acylation and deprotected thereafter. Other compounds of the invention may be prepared by reaction of a compound of formula (V) with a chloride of formula (VI) or an isocyanate of formula (VII):



The Examples herein provide further details of routes and methods for the preparation of compounds of the invention.

Compositions with which the invention is concerned may be prepared for administration by any route consistent with the pharmacokinetic properties of the active ingredient(s).

Orally administrable compositions may be in the form of tablets, capsules, powders, granules, lozenges, liquid or gel preparations, such as oral, topical, or sterile parenteral solutions or suspensions. Tablets and capsules for oral administration may be in unit dose presentation form, and may contain conventional excipients such as binding agents, for example syrup, acacia, gelatin, sorbitol, tragacanth, or polyvinyl-pyrrolidone; fillers for example lactose, sugar, maize-starch, calcium phosphate, sorbitol or glycine; tableting lubricant, for example magnesium stearate, talc, polyethylene glycol or silica; disintegrants for example potato starch, or acceptable wetting agents such as sodium lauryl sulphate. The tablets may be coated according to methods well known in normal pharmaceutical practice. Oral liquid preparations may be in the form of, for example, aqueous or oily suspensions, solutions, emulsions, syrups or elixirs, or may be presented as a dry product for reconstitution with water or other suitable vehicle before use. Such liquid preparations may contain conventional additives such as suspending agents, for example sorbitol, syrup, methyl cellulose, glucose syrup, gelatin hydrogenated edible fats; emulsifying agents, for example lecithin, sorbitan monooleate, or acacia; non-aqueous vehicles (which may include edible oils), for example almond oil, fractionated coconut oil, oily esters such as glycerine, propylene glycol, or ethyl alcohol; preservatives, for example methyl or propyl p-hydroxybenzoate or sorbic acid, and if desired conventional flavouring or colouring agents.

Safe and effective dosages for different classes of patient and for different disease states will be determined by clinical trial as is required in the art. It will be understood

that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following examples illustrate embodiments of the invention.:

In the Examples, the following abbreviations have been used:


DMF	Dimethylformamide
HOBt	1-Hydroxybenzotriazole
HPLC	High performance liquid chromatography
LRMS	Low resolution mass spectrometry
NMR	Nuclear magnetic resonance
RT	Retention Time
TLC	Thin layer chromatography
DIEA	N,N-diisopropylethylamine
DCM	Dichloromethane
HATU	O-(7-Azabenzotriazo-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate

^1H and ^{13}C NMR spectra were recorded using a Bruker DPX 250 spectrometer at 250.1 and 62.9MHz, respectively. Mass spectra were obtained using a Perkin Elmer Sciex API 165 spectrometer using both positive and negative ionisation modes. Infra-red spectra were recorded on a Perkin Elmer PE 1600 FTIR spectrometer.

Analytical HPLC was performed on a Beckman System Gold, using Waters Nova Pak C18 column (150 mm, 3.9 mm) with 20 to 90 % solvent B gradient (1 ml/min) as the mobile phase. [Solvent A: 0.05% TFA in 10% water 90% methanol; Solvent B: 0.05% TFA in 10% methanol 90%], detection wavelength at 230 nm. Preparative HPLC was performed on a Gilson autoprep instrument using a C18 Waters delta prep-pak cartridge (15 μm , 300 A, 25 mm, 10 mm) with 20 to 90 % solvent B gradient

(6 ml/min) as the mobile phase. [Solvent A water; Solvent B: methanol], UV detection was at 230 nm.

Example 1



Step A. 3-Benzyloxyformylamino-(2*R*)-cyclopentylmethyl-propionic acid, EDAC, HOBT, DMF, 0°C to rt, 5 h; Step B. 2-furoyl chloride, DIEA, CH₂Cl₂, 0°C to rt, 16 h; Step C. H₂ (g), 10% Pd/C, MeOH, rt, 2 h.

Step A:

N-Benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-(tetrahydropyridazin-1-yl)propyl]formamide

To a solution of 3-benzyloxyformylamino-(2*R*)-cyclopentylmethyl-propionic acid (3.73 g, 12.2 mmol) in DMF (40 ml) at 0°C under argon was added EDAC (2.58 g, 13.5 mmol) and HOBT (0.83 g, 6.12 mmol). The resulting mixture was stirred for 15 min at 0°C before 1,2-diazacyclohexane hydrochloride (1.50 g, 12.2 mmol) was introduced followed by DIEA (4.69 ml, 27.0 mmol) in DMF (5 ml). The reaction was maintained at 0°C for 1 h and then warmed to rt where it was left for 5 h. After this time, the reaction was diluted with EtOAc and washed with 1M Na₂CO₃. The aqueous phase was removed and extracted with additional EtOAc (x1). The combined organic fractions were washed with H₂O and sat. NaCl and then separated, dried (MgSO₄), filtered and concentrated under reduced pressure. The crude material was purified by chromatography on silica eluting with 100% EtOAc. This gave the title compound as a colourless oil (2.95 g, 65%). ¹H-NMR; δ (CD₃OD), 8.18 (0.5H, s, CHO), 7.88

(0.5H, s, CHO), 7.50-7.30 (5H, m, ArH), 4.92-4.65 (2H, br m, CH₂Ph), 4.10-2.80 (7H, m), 1.75-1.40 (13H, m), 1.20-1.00 (2H, m). LRMS: +ve ion: 396 [M+23], 374 [M+1], 346.

Step B:

N-Benzyloxy-*N*-{(2*R*)-cyclopentylmethyl-3-[2-(furan-2-carbonyl)-tetrahydropyridazin-1-yl]-3-oxo-propyl}formamide

To a solution of *N*-benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-(tetrahydropyridazin-1-yl)propyl]formamide (60 mg, 0.16 mmol) in DCM (1 ml) at 0°C under argon was added 2-furoyl chloride (17 µl, 0.17 mmol) followed by DIEA (34 µl, 0.19 mmol). The mixture was stirred for 30 min at 0°C and then warmed to rt where it was maintained for 16 h. After this time the reaction was diluted with EtOAc and washed with 1M citric acid, sat. NaHCO₃ and sat. NaCl. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure. The crude material was purified by chromatography on silica eluting with 70-100% EtOAc/hexanes. This gave the title compound as a colourless oil (67 mg, 89%). ¹H-NMR; δ (CD₃OD), 8.26 (0.33H, s, CHO), 7.95 (0.67H, s, CHO), 7.78 (1H, br s, HetArH), 7.50-7.25 (5H, m, ArH), 7.03 (1H, br m, HetArH), 6.63-6.52 (1H, br m, HetArH), 5.03-4.70 (2H, br m, CH₂Ph), 4.70-4.40 (2H, m), 4.30-2.85 (5H, m), 1.90-1.00 (13H, m), 0.95-0.60 (2H, m). LRMS: +ve ion: 490 [M+23].


Step C:

(2*R*)-Cyclopentylmethyl-3-[2-(furan-2-carbonyl)-tetrahydropyridazin-1-yl]-3-oxo-propyl}formamide

To a solution of the *N*-benzyloxy-*N*-{(2*R*)-cyclopentylmethyl-3-[2-(furan-2-carbonyl)-tetrahydropyridazin-1-yl]-3-oxo-propyl}formamide (65 mg, 0.14 mmol) in MeOH (3 ml) at rt under argon was added 10% Pd/C (8 mg). Hydrogen gas was bubbled through the black suspension for 15 min and then the reaction was left under an atmosphere of hydrogen for 2 h. After this time, the suspension was filtered and the collected solids were washed with additional MeOH. The combined filtrate and washings were concentrated under reduced pressure to give the title compound as a colourless solid (52 mg, 100%). ¹H-NMR; δ (CD₃OD), 8.29 (0.33H, s, CHO), 7.89

(0.67H, s, CHO), 7.78 (1H, br s, HetArH), 7.20-7.00 (1H, br m, HetArH), 6.64 (1H, br m, HetArH), 4.70-4.55 (2H, m), 3.95-2.95 (5H, m), 1.95-1.0 (13H, m), 0.95-0.80 (2H, m). LRMS: +ve ion: 400 [M+23]. HPLC Rt = 4.49 min (214 nm).

Example 2



Step A. 3,5-Bis-(trifluoromethyl)phenyl isocyanate, EtOAc, rt, 16 h; Step B. H₂ (g), 10% Pd/C, MeOH, rt, 2 h.

Step A:

2-[3-(Benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-tetrahydropyridazine-1-carboxylic acid (3,5-bis-(trifluoromethyl)phenyl)amide

To a solution of *N*-benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-(tetrahydropyridazin-1-yl)propyl]formamide (53 mg, 0.14 mmol) in EtOAc (2 ml) at rt under argon was added 3,5-bis-(trifluoromethyl)phenyl isocyanate (36 mg, 0.14 mmol). The resulting mixture was stirred for 16 h at rt and then concentrated under reduced pressure. The crude product was purified by chromatography on silica eluting with 70-100% EtOAc/hexanes to give the title compound as a colourless oil (76 mg, 85%). ¹H-NMR; δ(CD₃OD), 8.25-7.88 (3H, m, ArH and CHO), 8.60-7.20 (6H, m, ArH), 5.01-4.75 (2H, m, CH₂Ph), 4.60-2.90 (7H, m), 1.80-1.05 (13H, m), 1.00-0.90 (2H, m). LRMS: +ve ion: 651 [M+23]; 400.

Step B:


2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (3,5-bis-(trifluoromethyl)phenyl)amide

To a solution of the 2-[3-(benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-

tetrahydropyridazine-1-carboxylic acid (3,5-bis-(trifluoromethyl) phenyl)amide (75 mg, 0.12 mmol) in MeOH (4 ml) at rt under argon was added 10% Pd/C (10 mg). Hydrogen gas was bubbled through the black suspension for 15 min and then the reaction was left under an atmosphere of hydrogen for 2 h. After this time, the suspension was filtered and the collected solids were washed with additional MeOH. The combined filtrate and washings were concentrated under reduced pressure to give the title compound as a colourless solid (63 mg, 98%). ¹H-NMR; δ(CD₃OD), 8.30-8.24 (2.5H, m, ArH and CHO), 7.91 (0.5H, s, CHO), 7.62 (1H, s, ArH), 4.60-4.38 (2H, m), 4.10-2.85 (5H, m), 1.90-1.05 (13H, m), 1.00-0.90 (2H, m). LRMS: +ve ion: 561 [M+23]. HPLC Rt = 6.30 min (214 nm).

The compounds of Examples 3 to 9 were prepared by methods analogous to that of Example 2 substituting the appropriate isocyanate for that used in Example 2


Example 3



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid phenylamide

LRMS: +ve ion: 425 [M+23]. HPLC Rt = 5.13 min (214 nm).


Example 4



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (3-methoxyphenyl)amide

LRMS: +ve ion: 455 [M+23]. HPLC Rt = 5.13 min (214 nm).


Example 5



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (4-fluorophenyl)amide

LRMS: +ve ion: 443 [M+23]. HPLC Rt = 5.22 min (214 nm).


Example 6



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (2,4-bis(fluoro)phenyl)amide

LRMS: +ve ion: 461 [M+23]. HPLC Rt = 5.15 min (214 nm).


Example 7



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (4-trifluoromethylphenyl)amide


LRMS: +ve ion: 493 [M+23]. HPLC Rt = 5.81 min (214 nm).

Example 8




2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (2,3,4a,8a-tetrahydrobenzo[1,4]dioxin-6-yl)amide
LRMS: +ve ion: 483 [M+23]. HPLC Rt = 4.96 min (214 nm).

Example 9



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid (2-methoxyphenyl)amide
LRMS: +ve ion: 455 [M+23]. HPLC Rt = 5.27 and 5.37 min [rotamers] (214 nm).

Example 10



Step A. 3-Benzyloxyformylamino-(2*R*)-cyclopentylmethyl-propionic acid, HATU, DIEA, DMF, 0°C to rt, 2.5 h; Step B. H₂ (g), 10% Pd/C, MeOH, rt, 2 h.

Step A:

2-[3-(Benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-pyrazolidine-1-carboxylic acid *tert*-butyl ester

To a solution of 3-benzyloxyformylamino-(2*R*)-cyclopentylmethyl-propionic acid (103 mg, 0.34 mmol) in DMF (2 ml) at 0°C under argon was added HATU (128 mg, 0.34 mmol) and DIEA (70 µl, 0.41 mmol). The resulting mixture was stirred for 10 min before a solution of *N-tert*-(butoxycarbonyl)-1,2-diazacyclopentane (58 mg, 0.34 mmol) in DMF (0.5 ml) was introduced. After 1 h at 0°C, the reaction was warmed to rt where it was maintained for 1.5 h. The mixture was then diluted with EtOAc and washed with 1M citric acid, 1M Na₂CO₃ and sat. NaCl. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure. The crude product was purified by chromatography on silica eluting with 50-70% EtOAc/hexanes to give the title compound as a colourless oil (46 mg, 30%). ¹H-NMR; δ(CDCl₃), 8.23 (0.25H, s, CHO), 8.08 (0.25H, s, CHO), 7.99 (0.25 H, s, CHO), 7.85 (0.25H, s, CHO), 7.45-7.26 (5H, m, ArH), 4.87-4.60 (2H, br m, CH₂Ph), 4.20-2.80 (7H, m), 2.10-1.40 (11H, m), 1.45 (9H, s, (CH₃)₃), 1.20-1.00 (2H, m). LRMS: +ve ion: 482 [M+23], 382.


Step B:

2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid *tert*-butyl ester

To a solution of 2-[3-(benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-pyrazolidine-1-carboxylic acid *tert*-butyl ester (46 mg, 0.10 mmol) in MeOH (4 ml) at rt under argon was added 10% Pd/C (8 mg). Hydrogen gas was bubbled through the black suspension for 15 min and then the reaction was left under an atmosphere of hydrogen for 2 h. After this time, the suspension was filtered and the collected solids were washed with additional MeOH. The combined filtrate and washings were concentrated under reduced pressure to give the title compound as a colourless solid (36 mg, 97%). ¹H-NMR; δ(CD₃OD), 8.28 (0.25H, s, CHO), 8.22 (0.25H, s, CHO), 7.93 (0.25H, s, CHO), 7.80 (0.25H, s, CHO), 4.20-3.79 (4H, m), 3.78-3.40 (1H, m), 3.39-3.02 (2H, m), 2.20-1.40 (11H, m), 1.51 (4.5H, s, (CH₃)₃), 1.50 (4.5H, s, (CH₃)₃), 1.20-1.01 (2H, m). LRMS: +ve ion: 392 [M+23], 292, 270. HPLC Rt = 5.23 min (214 nm).

The compound of Example 11 was prepared by a method analogous to that to Example 10:


Example 11



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-tetrahydropyridazine-1-carboxylic acid *tert*-butyl ester

¹H-NMR; δ (CD₃OD), 8.26 (0.25H, s, CHO), 7.91 (0.25H, s, CHO), 7.84 (0.5H, s, CHO), 4.44-4.40 (1H, m), 4.22-3.10 (4H, m), 3.10-2.70 (2H, m), 1.85-1.30 (13H, m), 1.51 (4.5H, s, (CH₃)₃), 1.50 (4.5H, s, (CH₃)₃), 1.20-1.00 (2H, m). LRMS: +ve ion: 406 [M+23], 306. HPLC Rt = 5.41 and 5.55 min [rotamers] (214 nm).

Example 12



Step A. BF₃·Et₂O, CH₂Cl₂:AcOH (5:1), 0°C 2 h; Step B. 2-furoyl chloride, DIEA, CH₂Cl₂, 0°C to rt, 1 h; Step C. H₂ (g), 10% Pd/C, MeOH, rt, 1 h.

Step A:

N-Benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-pyrazolidin-1-yl-propyl]formamide
 To a solution of 2-[3-(benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-pyrazolidine-1-carboxylic acid *tert*-butyl ester (477 mg, 1.04 mmol) in DCM (12.5 ml) at 0°C under argon was added glacial acetic acid (2.5 ml) followed by boron trifluoride etherate (0.77 ml, 6.24 mmol). The solution was stirred at 0°C for 2 h and then diluted with EtOAc. The mixture was washed with 1M Na₂CO₃ and sat. NaCl. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure to give the title compound which was used without further purification. ¹H-NMR; δ(CDCl₃), 8.19 (0.5H, s, CHO), 7.88 (0.5H, s, CHO), 7.47-7.26 (5H, m, ArH), 4.95-4.60 (2H, br m, CH₂Ph), 3.93-3.30 (5H, m), 3.20-2.80 (2H, m), 2.00-1.40 (12H, m), 1.15-1.00 (2H, m). LRMS: +ve ion: 382 [M+23], 332.

Step B:

N-Benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-[2-(furan-2-carbonyl)-pyrazolidin-1-yl]-3-oxo-propyl]formamide

To a solution of *N*-benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-pyrazolidin-1-yl-propyl]formamide (31 mg, 0.086 mmol) in DCM (2 ml) at 0°C under argon was added 2-furoyl chloride (9 µl, 0.095 mmol) followed by DIEA (18 µl, 0.104 mmol). The mixture was stirred for 30 min at 0°C and then warmed to rt where it was maintained for 1 h. After this time the reaction was diluted with EtOAc and washed with 1M citric acid, sat. NaHCO₃ and sat. NaCl. The organic layer was dried (MgSO₄), filtered and concentrated under reduced pressure. The crude material was purified by chromatography on silica eluting with 80% EtOAc/hexanes. This gave the title compound as a colourless oil (34 mg, 87%). ¹H-NMR; δ(CD₃OD), 8.32 (0.25H, s, CHO), 8.10 (0.25H, s, CHO), 7.96 (0.25H, s, CHO), 7.90 (0.25H, s, CHO), 7.80-7.74 (1H, br m, HetArH), 7.50-7.20 (6H, br m, ArH and HetArH), 6.60 (1H, br s, HetArH), 5.00-4.80 (2H, br m, CH₂Ph), 4.40-2.95 (7H, m), 2.20-1.40 (11H, m), 1.20-0.85 (2H, m). LRMS: +ve ion: 476 [M+23].


Step C:

(2*R*)-Cyclopentylmethyl-3-[2-(furan-2-carbonyl)-pyrazolidin-1-yl]-3-oxo-

propyl}formamide

To a solution of the *N*-benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-[2-(furan-2-carbonyl)-pyrazolidin-1-yl]-3-oxo-propyl}formamide (32 mg, 0.071 mmol) in MeOH (2 ml) at rt under argon was added 10% Pd/C (5 mg). Hydrogen gas was bubbled through the black suspension for 15 min and then the reaction was left under an atmosphere of hydrogen for 1 h. After this time, the suspension was filtered and the collected solids were washed with additional MeOH. The combined filtrate and washings were concentrated under reduced pressure to give the title compound as a colourless solid (23 mg, 90%). ¹H-NMR; δ(CD₃OD), 8.28 (0.25H, s, CHO), 8.22 (0.25H, s, CHO), 7.90-7.70 (1.5H, m, CHO and HetArH), 7.30-7.20 (1H, br m, HetArH), 6.65 (1H, br s, HetArH), 4.40-3.05 (7H, m), 2.25-1.30 (11H, m), 1.20-0.85 (2H, m). LRMS: +ve ion: 386 [M+23]. HPLC Rt = 4.36 min (214 nm).

Example 13



Step A. 2-Methoxyphenyl isocyanate, EtOAc, rt, 16 h; Step B. H₂ (g), 10% Pd/C, MeOH, rt, 2 h.

Step A:

2-[3-(Benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-pyrazolidine-1-carboxylic acid (2-methoxyphenyl)amide

To a solution of *N*-benzyloxy-*N*-[(2*R*)-cyclopentylmethyl-3-oxo-3-pyrazolidin-1-yl-propyl]formamide (37 mg, 0.10 mmol) in EtOAc (2 ml) at rt under argon was added 2-methoxyphenyl isocyanate (15 mg, 0.10 mmol). The resulting mixture was stirred for 16 h at rt and then concentrated under reduced pressure. The crude product was purified by preparative HPLC to give the title compound as a colourless oil (28 mg,

53%). ¹H-NMR; δ(CD₃OD), 8.29 (0.5H, s, CHO), 8.04-7.90 (1.5H, CHO and ArH), 7.50-7.20 (5H, m, ArH), 7.23-6.87 (3H, ArH), 4.90-4.60 (2H, br m, CH₂Ph), 4.25-3.95 (7H, m), 3.83 (3H, s, CH₃O), 2.18-1.95 (2H, m), 1.90-0.85 (11H, m). LRMS: +ve ion: 531 [M+23], 360.


Step B:

2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid (2-methoxyphenyl)amide

To a solution of the 2-[3-(benzyloxyformylamino)-(2*R*)-cyclopentylmethyl-propionyl]-pyrazolidine-1-carboxylic acid (2-methoxyphenyl)amide (28 mg, 0.055 mmol) in MeOH (2 ml) at rt under argon was added 10% Pd/C (3 mg). Hydrogen gas was bubbled through the black suspension for 15 min and then the reaction was left under an atmosphere of hydrogen for 1 h. After this time, the suspension was filtered and the collected solids were washed with additional MeOH. The combined filtrate and washings were concentrated under reduced pressure to give the title compound as a colourless solid (18 mg, 78%). ¹H-NMR; δ(CD₃OD), 8.20-7.65 (2H, br m, CHO and ArH), 7.23-6.87 (3H, m, ArH), 4.30-3.10 (7H, m), 3.85 (1.5H, s, CH₃O), 3.84 (1.5H, s, CH₃O), 2.15-2.00 (2H, m), 1.90-0.90 (11H, m). LRMS: +ve ion: 441 [M+23]. HPLC Rt = 5.12 min (214 nm).

The compounds of Example 14-17 were prepared by methods analogous to that of Example 13


Example 14



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid phenylamide

LRMS: +ve ion: 411 [M+23]. HPLC Rt = 4.98 min (214 nm).


Example 15



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid (4-fluorophenyl)amide

LRMS: +ve ion: 429 [M+23]. HPLC Rt = 5.14 min (214 nm).


Example 16



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid (3-methoxyphenyl)amide

LRMS: +ve ion: 441 [M+23]. HPLC Rt = 5.04 min (214 nm).

Example 17



2-[(2*R*)-Cyclopentylmethyl-3-(formylhydroxyamino)-propionyl]-pyrazolidine-1-carboxylic acid (4-trifluoromethylphenyl)amide

LRMS: +ve ion: 479 [M+23]. HPLC Rt = 5.68 min (214 nm).

Biological Example


Minimal Inhibitory concentrations (MIC) of inhibitors against clinical isolates of *S. pneumoniae*, *H. influenzae* and *M. catarrhalis* obtained from the Public Health and Clinical Microbiology Laboratory, Addenbrooke's Hospital, Hills Road, Cambridge

CB2 2QW, UK, were determined by a standard agar plate dilution method following recommendations in **British Society for Antimicrobial Chemotherapy Working Party**. 1991. "A guide to sensitivity testing British Society for Antimicrobial Chemotherapy, London, United Kingdom". Briefly Iso-Sensitest agar (pH 7.2: Oxoid, United Kingdom) was employed supplemented with 5% horse blood (Oxoid) and 20 µg of NAD (Sigma) per ml to support growth of fastidious bacteria. The inoculum used was approximately 10^4 colony forming units of each isolate contained in a volume of 1 µl. Plates were incubated 18 to 24 hr in air, or for fastidious bacteria an atmosphere enriched with 4-6% carbon dioxide at 35°C. The MIC was determined as the lowest concentration of an antimicrobial tested that inhibited growth of the inoculum, disregarding a single persisting colony or faint haze caused by the inoculation.

The compounds of the Examples were are antibacterially active in the above assays, For example, the compound of Example 12 has MICs of 2 µg/ml or less against 6 strains of *S. pneumoniae*, less than 1 µg/ml against 4 strains of *H. influenzae* and less than 0.5 µg/ml against 3 strains of *M. catarrhalis*.

Claims

1. A compound of formula (I) or a pharmaceutically or veterinarily acceptable salt, hydrate or solvate thereof



wherein

Q represents a radical of formula $-N(OH)CH(=O)$ or formula $-C(=O)NH(OH)$;

Y represents $-C(=O)-$, $-C(=S)-$, $-S(=O)-$, or $-SO_2-$;

R_1 represents hydrogen, C_1-C_6 alkyl or C_1-C_6 alkyl substituted by one or more halogen atoms, or, except when Q is a radical of formula $-N(OH)CH(=O)$, a hydroxy, C_1-C_6 alkoxy, C_1-C_6 alkenyloxy, halogen, amino, C_1-C_6 alkylamino, or di- (C_1-C_6 alkyl)amino group;

R_2 represents a substituted or unsubstituted C_1-C_6 alkyl, C_1-C_3 alkyl-O- C_1-C_3 alkyl, C_1-C_3 alkyl-S- C_1-C_3 alkyl, cycloalkyl(C_1-C_3 alkyl)-, aryl(C_1-C_3 alkyl)-, heterocyclyl(C_1-C_3 alkyl)-, or $R^1R^2N-C_1-C_3$ alkyl group wherein R^1 represents hydrogen or C_1-C_3 alkyl and R^2 represents C_1-C_3 alkyl, or R^1R^2N- represents a cyclic amino group;

R_3 and R_4 taken together with the nitrogen atoms to which they are respectively attached form a saturated heterocyclic ring of from 4 to 7 ring atoms, which may be fused to a second carbocyclic or heterocyclic ring, either of which rings may optionally be substituted; and

A represents a primary, secondary or tertiary amino group or a group $-R_5$, $-OR_5$, wherein R_5 is a substituted or unsubstituted C_1-C_6 alkyl, C_2-C_6 alkenyl, C_2-C_6 alkynyl,

cycloalkyl, aryl, heterocyclyl, C₁-C₃ alkyl-O-C₁-C₃ alkyl, C₁-C₃ alkyl-S-C₁-C₃ alkyl, cycloalkyl(C₁-C₃ alkyl)-, heterocyclic(C₁-C₃ alkyl, aryl(C₁-C₃ alkyl)-, or R¹R²N-C₁-C₃ alkyl group wherein R¹ represents hydrogen or C₁-C₃ alkyl and R² represents C₁-C₃ alkyl, or R¹R²N- represents a cyclic amino group.

2. A compound as claimed in claim 1 wherein Q is an N-formyl hydroxylamine group -N(OH)CH(=O).

3 A compound as claimed in claim 1 or claim 2 wherein -Y- is -C(=O)- or -SO₂-.

4. A compound as claimed in any of the preceding claims wherein R₁ is hydrogen.

5. A compound as claimed in any of the preceding claims wherein R₂ is

optionally substituted C₁-C₆ alkyl, C₃-C₆ alkenyl, C₃-C₆ alkynyl or cycloalkyl;

phenyl(C₁-C₆ alkyl)-, phenyl(C₃-C₆ alkenyl)- or phenyl(C₃-C₆ alkynyl)- optionally substituted in the phenyl ring;

cycloalkyl(C₁-C₆ alkyl)-, cycloalkyl(C₃-C₆ alkenyl)- or cycloalkyl(C₃-C₆ alkynyl)- optionally substituted in the cycloalkyl ring; or


CH₃(CH₂)_pO(CH₂)_q- or CH₃(CH₂)_pS(CH₂)_q-, wherein p is 0, 1, 2 or 3 and q is 1, 2 or 3.

6. A compound as claimed in any of claims 1 to 4 wherein R₂ is methyl, ethyl, n- or iso-propyl, n- or iso-butyl, n-pentyl, iso-pentyl, 3-methyl-but-1-yl, n-hexyl, n-heptyl, n-octyl, methylsulfanylethyl, ethylsulfanylmethyl, 2-methoxyethyl, 2-ethoxyethyl, 2-ethoxymethyl, 3-hydroxypropyl, allyl, 3-phenylprop-3-en-1-yl, prop-2-yn-1-yl, 3-phenylprop-2-yn-1-yl, 3-(2-chlorophenyl)prop-2-yn-1-yl, but-2-yn-1-yl, cyclopentyl, cyclohexyl, cyclopentylmethyl, cyclopentylethyl, cyclopentylpropyl,

cyclohexylmethyl, cyclohexylethyl, cyclohexylpropyl, furan-2-ylmethyl, furan-3-methyl, tetrahydrofuran-2-ylmethyl, tetrahydrofuran-2-ylmethyl, piperidinylmethyl, pyrid-2-ylmethyl, pyrid-3-ylmethyl, pyrid-4-ylmethyl, phenylpropyl, 4-chlorophenylpropyl, 4-methylphenylpropyl, 4-methoxyphenylpropyl, benzyl, 4-chlorobenzyl, 4-methylbenzyl, or 4-methoxybenzyl.


7. A compound as claimed in any of claims 1 to 4 wherein R_2 is (C_1-C_6) alkyl-, cycloalkylmethyl-, (C_1-C_3) alkyl-S- (C_1-C_3) alkyl-, or (C_1-C_3) alkyl-O- (C_1-C_3) alkyl-, especially n-propyl, n-butyl, n-pentyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl or cyclohexylethyl.

8. A compound as claimed in any of the preceding claims wherein the ring formed by R_3 and R_4 and the nitrogens to which they are attached is one of the following, any of which may be optionally substituted, and wherein R represents hydrogen or C_1-C_4 alkyl and any sulfur atom present as a ring member may be oxidised to -SO- or -SO₂-:



9. A compound as claimed in any of the preceding claims wherein A is a secondary amino group or a cyclic or non-cyclic tertiary amino group.

10. A compound as claimed in any of claims 1 to 8 wherein A is an amino group of formula -NR₆R₇ wherein R₆ and R₇ independently represent a radical of formula (II)



wherein

m, p and n are independently 0 or 1;

Z represents hydrogen or a carbocyclic or heterocyclic ring of 5 to 7 ring atoms which is optionally fused to a saturated or unsaturated carbocyclic or heterocyclic second ring of 5 to 7 ring atoms;

Alk¹ and Alk² independently represent divalent C₁-C₃ alkylene radicals;

X represents -O-, -S-, -S(O)-, -S(O₂)-, -C(=O)-, -NH-, -NR₇- where R₇ is C₁-C₃ alkyl;

and wherein

Alk¹, Alk² and Z when other than hydrogen, independently are optionally substituted by

(C₁-C₃)alkyl, (C₂-C₃)alkenyl, or (C₂-C₃)alkynyl,
phenyl, optionally substituted by (C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo, nitro,
amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl;

monocyclic 5 or 6-membered heterocyclic, optionally substituted by (C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo, nitro, amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl

benzyl, optionally substituted in the phenyl ring by (C₁-C₃)alkyl, (C₁-C₃)alkoxy, halo, nitro, amino, mono- or di-(C₁-C₃)alkylamino, cyano or trifluoromethyl,

hydroxy, phenoxy, (C₁-C₆)alkoxy, or hydroxy(C₁-C₆)alkyl,

mercapto, (C₁-C₆)alkylthio or mercapto(C₁-C₆)alkyl,

oxo,

nitro,

cyano

halo

-COOH, or -COOR^A,

-CONH₂, -CONHR^A, or -CONR^AR^B

-COR^A, -SO₂R^A,

-NHCOR^A,

-NH₂, -NHR^A, or -NR^AR^B,

wherein R^A and R^B are independently a (C₁-C₆) alkyl group, R^A and R^B taken together with the nitrogen atom to which they are attached form a 5- or 6-membered heterocyclic ring which may be substituted by (C₁-C₃)alkyl, hydroxy, or hydroxy(C₁-C₃)alkyl.

11. A compound as claimed in claim 10 wherein Alk¹ and Alk² independently represent -(CH₂)- or -(CH₂CH₂)-.

12. A compound as claimed in claim 10 or claim 11 wherein m is 0, p is 1, n is 0 or 1 and X is -C(=O)- or -S(O₂)-.



13. A compound as claimed in claim 10 wherein the substituent (II) has the formula -CH₂Z, -OZ, or -(C=O)Z wherein Z is C₁-C₃ alkyl, phenyl, 3,4-methylenedioxyphenyl, morpholinyl, pyrimidinyl, 1,2,3-thiadiazolyl, 1,4-thiazolyl, benzofuranyl, furanyl, thienyl, pyranyl, pyrrolyl, pyrazolyl, isoxazolyl, or pyridyl, any of which may optionally be substituted as specified. In particular, Z may be a methyl, ethyl, n- or iso-propyl, phenyl, 3,4-methylenedioxyphenyl, morpholinyl, pyrimidin-2-yl, 1,2,3-thiadiazol-5-yl, 1,4-thiazol-5-yl, benzofuran-2-yl, 2- or 3-furanyl, 2- or 3-thienyl, 2- or 3-pyranyl, 2-, 3- or 4-pyrrolyl, 3-, 4- or 5-pyrazolyl, 3-, 4- or 5-isoxazolyl, or 2-, 3- or 4-pyridyl ring any of which may optionally be substituted as specified in the broad description of the compounds of the invention.

14. A compound as claimed in any of claims 1 to 8 wherein A is an amino group of formula -NR₆R₇ wherein R₆ and R₇ taken together with the nitrogen atom to which they are attached form a saturated heterocyclic ring of 5 to 8 atoms optionally fused to a saturated or unsaturated carbocyclic or heterocyclic second ring of 5 to 7 ring

atoms, any of which rings being optionally substituted by a radical of formula (II) as defined in any of claims 10 to 13.

15. A compound as claimed in claim 14 wherein A is optionally substituted piperidin-1-yl or 1-piperazinyl.

16. A compound as claimed in claim 1 of formula (IIA) or (IIB)



wherein R_2 is as defined in claim 1;

X_1 is a bond, C_1 - C_3 alkylene, -NH- or -O-; and

A_1 is optionally substituted C_1 - C_6 alkyl, cycloalkyl, aryl, or heterocyclic.

17. A compound as claimed in claim 16 wherein R_2 is n-propyl, n-butyl, n-pentyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl or cyclohexylethyl;

X_1 is a bond, - CH_2 -, - CH_2CH_2 -, - $CH_2CH_2CH_2$ -, -NH- or -O-; and

A_1 is methyl, ethyl phenyl, cyclopentyl, cyclohexyl, 2- or 3-furanyl, 2- or 3-thienyl, 2-, 3- or 4-pyridyl, 3-, 4- or 5-pyrazolyl, 3-, 4- or 5-oxazolyl, or 3-, 4- or 5-thiazolyl, methoxymethyl, 3,5-bis-(trifluoromethyl)phenyl, 4-trifluoromethylphenyl, 4-methoxyphenyl, 3,4-methylenedioxyphenyl, 4-fluorophenyl benzyl, 3-pyridyl, 4-pyridyl, cyclohexyl, 1,3-dimethylpyrazol-5-yl, 1-methylimidazol-5-yl, or 2-[morpholin-1-yl]pyrid-5-yl.

18. A method for the treatment of bacterial infections in humans and non-human mammals, which comprises administering to a subject suffering such infection an antibacterially effective dose of a compound as claimed in any of the preceding claims.
19. The use of a compound as claimed in any of claims 1 to 17 for inhibiting bacterial growth in vitro and in vivo in mammals.
20. The use of a compound as claimed in any of claims 1 to 17 for the manufacture of a composition for treating bacterial infection by inhibiting bacterial growth.
21. A method for the treatment of bacterial contamination by applying an antibacterially effective amount of a compound as claimed in any of claims 1 to 17 to the site of contamination.
22. A pharmaceutical or veterinary composition comprising a compound as claimed in any of claims 1 to 17 together with a pharmaceutically or veterinarily acceptable carrier or excipient.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.